



# Marine Virus-Like Particles and Microbes: A Linear Interpretation

B. B. Cael<sup>1,2\*</sup>, Michael C. G. Carlson<sup>3</sup>, Christopher L. Follett<sup>1</sup> and Michael J. Follows<sup>1</sup>

<sup>1</sup> Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, United States, <sup>2</sup> Woods Hole Oceanographic Institution, Woods Hole, MA, United States, <sup>3</sup> Faculty of Biology, Technion, Israel Institute of Technology, Haifa, Israel

**Keywords:** marine viruses, virus, viroplankton, viral abundance, virus-like particles, virus to microbe ratio, virus to bacterium ratio, virus-microbe relationship

Viruses are key players in ocean ecology and biogeochemistry, not only because of their functional roles but also partially due to their sheer abundance (Fuhrman, 1999; Wilhelm and Suttle, 1999). Because viruses cannot replicate without their hosts' machinery, their abundance is inextricably related to that of their (mostly microbial) hosts. The relationship between viral and microbial abundances is thus of great interest.

It is often assumed that the abundance of virus-like particles ( $V$ ,  $\text{mL}^{-1}$ ) and microbial cells ( $M$ ,  $\text{mL}^{-1}$ ) are approximately proportional. A rule-of-thumb "virus-to-microbe" ratio of 10 has been used (Thingstad, 2000) as measurements of  $V$  are frequently an order of magnitude larger than those of  $M$ . However,  $V/M$  is known to vary substantially (Maranger and Bird, 1995; Knowles et al., 2016; Wigington et al., 2016; Parikka et al., 2017).

Wigington et al. (2016) presented an alternative paradigm. They compiled 5,671  $V$  and  $M$  measurements from 25 studies spanning diverse marine environments, an order of magnitude larger than any previous data compilation of its kind. They definitively showed that a power-law relationship  $V = \alpha M^\beta$  is a better statistical model than a proportionality  $V = \alpha M$ , in terms of the proportion of variance explained by each. They found  $\beta < 1$  in most cases, meaning  $V/M$  decreases as  $M$  increases. If  $V$  does scale nonlinearly with  $M$  this has broad implications for the characterization of marine viruses' abundance and influence (Wigington et al., 2016), so it is important to be certain that a nonlinear model is superior to a linear description. It also raises questions as to how this nonlinear scaling emerges, and what determines  $\alpha$  and  $\beta$ .

While several studies (Maranger and Bird, 1995; Danovaro et al., 2011; Knowles et al., 2016; Wigington et al., 2016) have tested the model  $V = \alpha M^\beta$  by performing ordinary least-squares (OLS) regression on log-transformed  $V$  and  $M$  data, to our knowledge none have tested the linear model  $V = aM + b$ . Unlike  $V = \alpha M$ , the linear model  $V = aM + b$  has the same number of parameters as the nonlinear model  $V = \alpha M^\beta$ , is consistent with a decreasing  $V/M$  as  $M$  increases, and is not a special case of the nonlinear model. We interpret  $V = aM + b$  to consider  $V$  as the sum of two pools—a pool of viruses whose abundance is proportional to microbial abundance ( $aM$ ), and a background pool of other viruses and virus-like particles whose abundance is unrelated to microbial abundance ( $b$ ). Because this model is simple and its parameters are comparatively straightforward to interpret and possibly predict, its ability to explain observations should be evaluated before concluding that  $V$  and  $M$  are nonlinearly related.

Here we evaluate the ability of the linear model  $V = aM + b$  to explain the relationship between  $V$  and  $M$  in the marine datasets considered by Wigington et al. (2016), Knowles et al. (2016), and Parikka et al. (2017). We show that the linear model's performance is never significantly different from that of the nonlinear model, and therefore that the linear model is a tenable description of the relationship between  $V$  and  $M$ .

We first reanalyzed the datasets considered in Wigington et al. (2016). They analyzed data from 25 studies, and also aggregated surface ( $z \leq 100$  m;  $n = 2,921$ ) and subsurface ( $z > 100$  m;  $n = 2,750$ ) samples, totaling 27 datasets. We repeated their OLS regression of log<sub>10</sub>-transformed  $V$  and  $M$  data (see Supplementary Material for MATLAB R2017a code). We also fit the model  $V = aM + b$  after log-transformation:

## OPEN ACCESS

### Edited by:

Ian Hewson,  
Cornell University, United States

### Reviewed by:

Stéphan Jacquet,  
Institut National de la Recherche  
Agronomique (INRA), France  
Gian Marco Luna,  
Consiglio Nazionale Delle Ricerche  
(CNR), Italy

### \*Correspondence:

B. B. Cael  
snail@mit.edu

### Specialty section:

This article was submitted to  
Aquatic Microbiology,  
a section of the journal  
Frontiers in Microbiology

Received: 14 November 2017

Accepted: 15 February 2018

Published: 01 March 2018

### Citation:

Cael BB, Carlson MCG, Follett CL and  
Follows MJ (2018) Marine Virus-Like  
Particles and Microbes: A Linear  
Interpretation. *Front. Microbiol.* 9:358.  
doi: 10.3389/fmicb.2018.00358



as a description of the relationship between viral and microbial abundances.

The linear model can be interpreted as decomposing virus-like particle abundance into two terms:  $V = aM + b = V_a + V_b$ . In this interpretation, the first term  $V_a$  is the abundance of viruses that infect organisms included in  $M$ , and it is assumed that  $V_a \propto M$  as captured by the parameter  $a$ . In marine systems,  $M$  is dominated by the abundance of prokaryotes, thus  $V_a$  can be considered to generally represent the abundance of bacteriophage (Suttle, 2007).  $a$  can then be considered a characteristic virus-to-microbe ratio, and can be understood in the context of dynamical models. For example, consider arguably the simplest model for virus-microbe dynamics (Lotka, 1920; Volterra, 1928; Lauro et al., 2011; Yau et al., 2011),

$$\frac{dV}{dt} = \gamma\varphi VM - \lambda V, \quad \frac{dM}{dt} = \mu M - \varphi VM$$

where ( $\gamma$ ,  $\varphi$  [mL s<sup>-1</sup>],  $\lambda$  [s<sup>-1</sup>],  $\mu$  [s<sup>-1</sup>]) represent burst size, virus-microbe interactions, viral decay, and microbial growth respectively (Record et al., 2016). At steady state, this model predicts  $V/M = \gamma\mu/\lambda$ . Predictions for  $a$  can therefore be derived from such models. In the above model,  $a$  is the product of the viral burst size, the host growth rate, and the viral decay timescale—though most of these are poorly parameterized in environmental populations. Across the 29 datasets we considered, our estimates of  $a$  ranged from 0 to 92, with a median of 8.2 (Table S1). This large variation is similar to that of  $\alpha$  and  $\beta$  (Table S1; Knowles et al., 2016; Wigington et al., 2016), and is also consistent with the large range in virus-to-microbe ratios observed across marine environments (Parikka et al., 2017). In a majority of cases  $a$  was within a factor of two of the “rule of thumb” virus-to-microbe ratio of 10.

The second term  $V_b$  is an additional “background pool” of virus-like particles, whose abundance is assumed to be unrelated to microbial abundance. Different types of particles other than infective bacteriophage may be present in this background pool, which could plausibly be variable and non-negligible. Virus-like particles are operationally defined as particles < 0.2  $\mu\text{m}$  in size containing nucleic acids; their abundance is measured by treating samples with a fluorescent nucleic acid stain, then counting fluorescing particles with epifluorescence microscopy (Noble and Fuhrman, 1998; Patel et al., 2007). Several non-viral sources are known to be captured by these methods whose contribution to total counts can be large, such as free nucleic acids (Bettarel et al., 2000), DNA-containing extracellular vesicles that are secreted by numerous marine microbes (Soler et al., 2015; Biller et al., 2017), gene-transfer agents (Biers et al., 2008), and decomposing viral material (Wommack et al., 1996). Additionally, while the relationship between microbes and viruses is generally thought to be dominated by prokaryotes and bacteriophage, eukaryote-infecting viruses (Nagasaki and Bratbak, 2010), zooplankton-infecting viruses (Fischer et al., 2010), and viroplage (La Scola et al., 2008) may also contribute to  $V$  without their corresponding hosts contributing to  $M$ . Finally, virus-like particles and microbes have overlapping sizes and counting each population has a significant degree of subjectivity. Therefore, many different

kinds of particles might be subsumed into  $V_b$ ; this background pool could change across environmental gradients (Biller et al., 2017) and with host types in ways unrelated to microbial density. Predictions for  $b$  could be made depending on which combination of these particle types  $b$  is assumed to represent. Across the 29 datasets we considered, our estimates of  $b$  also ranged substantially, from 0 to  $6 \times 10^7$  mL<sup>-1</sup>, with a median of  $6 \times 10^5$  mL<sup>-1</sup>. Biller et al. (2014) reported vesicle abundances of  $3 \times 10^5$  and  $6 \times 10^6$  mL<sup>-1</sup> for coastal surface water and Sargasso Sea samples; interestingly, a majority of our estimates for  $b$  were between these two values (Table S1).

Interpreting the nonlinear model is less straightforward. Knowles et al. (2016) developed a theory of viruses switching from lytic to lysogenic lifestyles to explain the sublinear scaling relationships they observed, but their theory has been disputed (Weitz et al., 2017) and broadly speaking the relationships between viral and microbial abundances they find tend to be weak or have  $\beta \approx 1$  (not indicative of nonlinearity); see their Figures 1a, 2. Empirical sublinear relationships are ubiquitous in biology (Hatton et al., 2015), but the underpinnings of the metabolic theory commonly invoked to explain these relationships have also been disputed (Dodds et al., 2001). It is unclear how the nonlinear model would account for virus-like particles other than bacteriophage, or how  $\alpha$  and  $\beta$  would be interpreted and predicted.

These analyses highlight the need for alternative methods to quantify viruses in the environment relative to their hosts. Measurements from marine environments with extremely low microbial densities (because the two models diverge for low values of  $M$ ), novel experiments with model systems, or other types of data that improve upon the indiscriminant measure of virus-like particles may help answer this question (Baran et al., 2018), and may also be useful for making accurate estimates of  $b$ . For now, we contest that a linear model remains plausible.

## AUTHOR CONTRIBUTIONS

BC conceived the research, performed the analyses, and wrote the paper. MC, CF, and MF assisted in the writing process and revised the manuscript.

## FUNDING

The National Science Foundation (GRFP-2388357, OCE-1315201), the National Aeronautics and Space Administration (NNX16AR47G), and the Simons Foundation (SCOPE Award 329108, CBIOMES Award 553242) supported this work.

## ACKNOWLEDGMENTS

We thank Maureen L. Coleman for productive discussions and suggestions.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2018.00358/full#supplementary-material>

## REFERENCES

- Baran, N., Goldin, S., Maidanik, I., and Lindell, D. (2018). Quantification of diverse virus populations in the environment using the polony method. *Nat. Microbiol.* 3, 62–72. doi: 10.1038/s41564-017-0045-y
- Bettarel, Y., Sime-Ngando, T., Amblard, C., and Laveran, H. (2000). A comparison of methods for counting viruses in aquatic systems. *Appl. Environ. Microbiol.* 66, 2283–2289. doi: 10.1128/AEM.66.6.2283-2289.2000
- Biers, E. J., Wang, K., Pennington, C., Belas, R., Chen, F., and Moran, M. A. (2008). Occurrence and expression of gene transfer agent genes in marine bacterioplankton. *Appl. Environ. Microbiol.* 74, 2933–2939. doi: 10.1128/AEM.02129-07
- Billler, S. J., Schubotz, F., Roggensack, S. E., Thompson, A. W., Summons, R. E., and Chisholm, S. W. (2014). Bacterial vesicles in marine ecosystems. *Science* 343, 183–186. doi: 10.1126/science.1243457
- Billler, S. J., McDaniel, L. D., Breitbart, M., Rogers, E., Paul, J. H., and Chisholm, S. W. (2017). Membrane vesicles in sea water: heterogeneous DNA content and implications for viral abundance estimates. *ISME J.* 11, 394–404. doi: 10.1038/ismej.2016.134
- Danovaro, R., Corinaldesi, C., Dell'Anno, A., Fuhrman, J. A., Middelburg, J. J., Noble, R. T., et al. (2011). Marine viruses and global climate change. *FEMS Microbiol. Rev.* 35, 993–1034. doi: 10.1111/j.1574-6976.2010.00258.x
- Dodds, P. S., Rothman, D. H., and Weitz, J. S. (2001). Re-examination of the “3/4-law” of metabolism. *J. Theor. Biol.* 209, 9–27. doi: 10.1006/jtbi.2000.2238
- Efron, B. (1979). Bootstrap methods: Another look at the jackknife. *Anna. Statist.* 7, 1–26. doi: 10.1214/aos/1176344552
- Fischer, M. G., Allen, M. J., Wilson, W. H., and Suttle, C. A. (2010). Giant virus with a remarkable complement of genes infects marine zooplankton. *Proc. Natl. Acad. Sci. U.S.A.* 107, 19508–19513. doi: 10.1073/pnas.1007615107
- Fuhrman, J. A. (1999). Marine viruses and their biogeochemical and ecological effects. *Nature* 399, 541–548. doi: 10.1038/21119
- Hatton, I. A., McCann, K. S., Fryxell, J. M., Davies, T. J., Smerlak, M., Sinclair, A. R., et al. (2015). The predator-prey power law: Biomass scaling across terrestrial and aquatic biomes. *Science* 349:aac6284. doi: 10.1126/science.aac6284
- Knowles, B., Silveira, C. B., Bailey, B. A., Barott, K., Cantu, V. A., Cobian-Guemes, A. G., et al. (2016). Lytic to temperate switching of viral communities. *Nature* 531, 466–470. doi: 10.1038/nature17193
- La Scola, B., Desnues, C., Pagnier, I., Robert, C., Barrassi, L., Fournous, G., et al. (2008). The virophage as a unique parasite of the giant mimivirus. *Nature* 455, 100–104. doi: 10.1038/nature07218
- Lauro, F. M., DeMaere, M. Z., Yau, S., Brown, M. V., Ng, C., Wilkins, D., et al. (2011). An integrative study of a meromictic lake ecosystem in Antarctica. *ISME J.* 5, 879–895. doi: 10.1038/ismej.2010.185
- Lotka, A. J. (1920). Analytical note on certain rhythmic relations in organic systems. *Proc. Natl. Acad. Sci. U.S.A.* 6, 410–415. doi: 10.1073/pnas.6.7.410
- Maranger, R., and Bird, D. F. (1995). Viral abundance in aquatic systems: a comparison between marine and fresh waters. *Marine Ecol. Progr. Ser.* 121, 217–226. doi: 10.3354/meps121217
- Muck, S., Griessler, T., Köstner, N., Klimiuk, A., Winter, C., and Herndl, G. J. (2014). Fracture zones in the Mid Atlantic Ridge lead to alterations in prokaryotic and viral parameters in deep-water masses. *Front. Microbiol.* 5:264. doi: 10.3389/fmicb.2014.00264
- Nagasaki, K., and Bratbak, G. (2010). “Isolation of viruses infecting photosynthetic and nonphotosynthetic protists,” in *Manual of Aquatic Viral Ecology*, eds.
- S. W. Wilhelm, M. G. Weinbauer, and C. A. Suttle (ASLO), 92–101. doi: 10.4319/mave.2010.978-0-9845591-0-7.92
- Noble, R. T., and Fuhrman, J. A. (1998). Use of SYBR Green I for rapid epifluorescence counts of marine viruses and bacteria. *Aquat. Microb. Ecol.* 14, 113–118. doi: 10.3354/ame014113
- Parikka, K. J., Le Romancer, M., Wauters, N., and Jacquet, S. (2017). Deciphering the virus-to-prokaryote ratio (VPR): insights into virus-host relationships in a variety of ecosystems. *Biol. Rev.* 92, 1081–1100. doi: 10.1111/brv.12271
- Parsons, R. J., Breitbart, M., Lomas, M. W., and Carlson, C. A. (2012). Ocean time-series reveals recurring seasonal patterns of virioplankton dynamics in the northwestern Sargasso Sea. *ISME J.* 6, 273–284. doi: 10.1038/ismej.2011.101
- Patel, A., Noble, R. T., Steele, J. A., Schwalbach, M. S., Hewson, I., and Fuhrman, J. A. (2007). Virus and prokaryote enumeration from planktonic aquatic environments by epifluorescence microscopy with SYBR Green I. *Nat. Protocols* 2, 269–276. doi: 10.1038/nprot.2007.6
- Record, N. R., Talmy, D., and Våge, S. (2016). Quantifying tradeoffs for marine viruses. *Front. Mar. Sci.* 3:251. doi: 10.3389/fmars.2016.00251
- Soler, N., Krupovic, M., Marguet, E., and Forterre, P. (2015). Membrane vesicles in natural environments: a major challenge in viral ecology. *ISME J.* 9:793. doi: 10.1038/ismej.2014.184
- Suttle, C. A. (2007). Marine viruses-major players in the global ecosystem. *Nat. Rev. Microbiol.* 5, 801–812. doi: 10.1038/nrmicro1750
- Thingstad, T. F. (2000). Elements of a theory for the mechanisms controlling abundance, diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems. *Limnol. Oceanogr.* 45, 1320–1328. doi: 10.4319/lo.2000.45.6.1320
- Volterra, V. (1928). Variations and fluctuations of the number of individuals in animal species living together. *ICES J. Marine Sci.* 3, 3–51.
- Weitz, J. S., Beckett, S. J., Brum, J. R., Cael, B. B., and Dushoff, J. (2017). Lysis, lysogeny and virus-microbe ratios. *Nature* 549, E1–E3. doi: 10.1038/nature23295
- Wigington, C. H., Sonderegger, D., Brussaard, C. P., Buchan, A., Finke, J. F., Fuhrman, J. A., et al. (2016). Re-examination of the relationship between marine virus and microbial cell abundances. *Nat. Microbiol.* 1:15024. doi: 10.1038/NMICROBIOL.2015.24
- Wilhelm, S. W., and Suttle, C. A. (1999). Viruses and nutrient cycles in the sea: viruses play critical roles in the structure and function of aquatic food webs. *Bioscience* 49, 781–788.
- Wommack, K. E., Hill, R. T., Muller, T. A., and Colwell, R. R. (1996). Effects of sunlight on bacteriophage viability and structure. *Appl. Environ. Microbiol.* 62, 1336–1341.
- Yau, S., Lauro, F. M., DeMaere, M. Z., Brown, M. V., Thomas, T., Raftery, M. J., et al. (2011). Virophage control of antarctic algal host-virus dynamics. *Proc. Natl. Acad. Sci. U.S.A.* 108, 6163–6168. doi: 10.1073/pnas.1018221108

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Cael, Carlson, Follett and Follows. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.